



For use by federal, state, tribal, or local government officials responsible for public health or vector control, or by persons certified in the appropriate category or otherwise authorized by the state or tribal lead pesticide regulatory agency to perform adult mosquito control applications, or by persons under their direct supervision.

This product is Ready-To-Use, Quick Knockdown, Low Odor, Synergized Synthetic Pyrethroid for Control of Adult Mosquitoes, Biting and Non-Biting Midges, Black flies and other outdoor flying insects as listed on this label which cause Public Health annoyance in Residential, Municipal and Recreational Areas.

READ ENTIRE LABEL BEFORE USING

READY-TO-USE

ACTIVE INGREDIENTS:	
*Permethrin.....	4.0%
**Piperonyl Butoxide.....	4.0%
OTHER INGREDIENTS.....	
	92.0%
TOTAL:	100.0%
*(3-phenoxyphenyl)methyl(+/-)Cis/trans 3-(2,2-dichloroethenyl)2,2-Dimethyl cyclopropanecarboxylate.cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.	
**(butylcarbityl)(6-propylpiperonyl) ether and related compounds.	
Contains 0.33 pounds of Permethrin and 0.33 pounds of Piperonyl Butoxide per gallon.	

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente" (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
If Swallowed:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Hot Line Number:	Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time. During other times, call the poison control center at 1-800-222-1222.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic organisms, including fish and aquatic invertebrates. Runoff from treated areas or deposition of spray droplets into a body of water may be hazardous to fish and aquatic invertebrates. Do not apply over bodies of water (lakes, rivers, permanent streams, natural ponds, commercial fish ponds, swamps, marshes or estuaries), except when necessary to target areas where adult mosquitoes are present, and weather conditions will facilitate movement of applied material away from the water in order to minimize incidental deposition into the water body. Do not contaminate bodies of water when disposing of equipment rinsate or washwaters.

BEE WARNING: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply to blooming crops or weeds while bees are visiting the treatment areas, except when applications are made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AllPro Envion rtu is recommended for application as a Non-Thermal aerosol (cold fog) to control adult mosquitoes and other insects as listed on this label in residential and recreational areas where these insect pests become a problem; such as, but not limited to parks, campsites, woodlands, athletic fields, golf courses, residential areas, municipalities, gardens, playgrounds, and non-crop waste areas.

This product is not for use in outdoor residential misting systems.

AllPro Envion rtu can be applied over specific growing crops and range grasses prior to harvest for the control of adult mosquitoes and biting flies within or adjacent to these areas. Application can only be made where the following crops are present: Alfalfa, Almonds, Apples, Artichokes (globe), Asparagus, Avocados, Broccoli, Brussels sprouts, Cabbage, Cauliflower, Celery, Cherries, Corn (forage), Corn (fodder), Corn (grain: field and pop), Corn (stover), Corn (sweet kernel + cob with husks removed), Eggplant, Filberts, Garlic, Kiwifruit, Horseradish, Leafy vegetables (except Brassica), Lettuce (head), Mushrooms, Onions (dry bulb), Peaches, Pears, Peppers (bell), Pistachios, Potato, Range grasses, Soybeans, Spinach, Tomatoes, Vegetables (cucurbit), Walnuts, Watercress

Do not allow spray treatment to drift on cropland (other than crops listed) or water supplies. In treatment of corrals, feedlots, animal confinements/houses, swine lots and zoos, cover any exposed drinking water, drinking fountains and animal feed before application.

WHEN TO APPLY

NOTE: Before making the first application in a season, it is advisable to consult with the state or tribal agency with primary responsibility for pesticide regulation to determine if other regulatory requirements exist.

FOR BEST RESULTS: Treat when mosquitoes or insects are most active and weather conditions are conducive to keeping the fog near the ground, e.g. cool temperatures, and consistent wind speeds are not greater than 10 MPH. Apply when wind speed is greater or equal to 1 MPH. Applications during the cool hours of night or early morning are usually preferable. Do not apply when air temperature is below 50° F.

Do not retreat the same site more than once in 12 hours. Do not exceed 25 applications at 0.007 pound permethrin A.I. per acre per application or 0.025 pound PBO A.I. per acre per application, or 0.18 pound of permethrin per acre per season or 2.0 pounds PBO per acre per year. More frequent treatments may be made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

HOW TO APPLY

GROUND APPLICATION: Spray equipment must be adjusted so the volume median diameter is less than 30 microns (Dv 0.5 < 30 µm) and that 90% of the spray is contained in droplets smaller than 50 microns (Dv 0.9 < 50 µm). Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

For proper ground application mount the fog applicator so that the nozzle is at least 4 1/2 ft. above the ground level and directed out the back of the vehicle. Failure to follow these label directions may result in reduced effectiveness.

AERIAL APPLICATION: Do not apply by fixed wing aircraft at a height less than 100 ft., or by helicopter at a height less than 75 ft. unless specifically approved by the state or tribe based on public health needs. The effects of flight speed and, for non-rotary nozzles, nozzle angle on the droplet size spectrum must be considered. Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a wind tunnel and laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

AERIAL APPLICATION MADE AT ≤ 200 FEET ABOVE GROUND ELEVATION: Spray equipment must be adjusted so that the volume median diameter produced is less than 60 microns (Dv 0.5 < 60 µm) and that 90% of the spray is contained in droplets smaller than 100 microns (Dv 0.9 < 100 µm).

AERIAL APPLICATION MADE AT > 200 FEET ABOVE GROUND ELEVATION: Spray equipment must be adjusted so the volume median diameter produced is less than 70 microns (Dv 0.5 < 70 µm) and that 90% of the spray is contained in droplets smaller than 145 microns (Dv 0.9 < 145 µm).

GROUND ULV NON-THERMAL AEROSOL (COLD FOG) APPLICATIONS

To control Mosquitoes, Midges and Black flies, apply **AllPro Envion rtu** using any standard ULV ground applicator capable of producing a non-thermal aerosol fog and base all acreage calculations on the equipment manufacturer's recommended swath width.

Apply at a flow rate of 2.35 to 16.46 ounces per minute at an average vehicle speed of 10 MPH. These rates are equivalent to 0.001 to 0.007 pound of permethrin A.I. and 0.001 to 0.007 pound of piperonyl butoxide A.I. per acre. If a different vehicle speed is used, adjust the rate accordingly. Vary flow rates accordingly to vegetation density and mosquito population. Use higher flow rates in heavy vegetation or when insect populations are high. An accurate flow rate control system must be used to ensure proper flow rate. Do not exceed a flow rate of 20 ounces per minute unless recommended by equipment manufacturers and/or Public Health Officials, high vehicle speeds of 10, 15 or 20 MPH are necessary in emergency conditions, and when fogging equipment is capable of such higher rates of output per minute.

USE AS AN INSECT BARRIER TREATMENT

For use in Non-Thermal ULV, motorized, portable backpack fog type mist blower equipment, apply **AllPro Envion rtu** with equipment adjusted to deliver ULV particles of 50 to 100 volume median diameter. Use 0.001 to 0.007 pound of permethrin A.I. per acre at a 50 ft. (15.2 m.) swath width while walking at a speed of 2 MPH (3.2 KPH). If applied at a different speed, make sure that the same amount (0.001 to 0.007 pound) of permethrin A.I. and piperonyl butoxide A.I. is applied per acre.

When this product is applied as a barrier application using low pressure hand sprayers, hydraulic sprayers or ground ULV equipment, do not apply within 100 ft. (30 m.) of lakes and streams.

AERIAL APPLICATION

Aerial application with ULTRA LOW VOLUMES of **AllPro Envion rtu** applied according to the rate charts on this label may be considered necessary when conducted at the discretion of Public Health Officials, Mosquito Abatement Districts, or other trained personnel engaged in outdoor Mosquito and flying insect control programs, when these insects threaten to become a public health hazard.

Aerial application of **AllPro Envion rtu** shall be made at altitudes ranging from 100 ft. to 300 ft. The recommended airspeed for Fixed Wing aircraft with flat fan nozzles is from 120 to 160 MPH. For Rotary Aircraft equipped with rotary nozzles, apply at a sufficient airspeed to deliver the appropriate amount of **AllPro Envion rtu** A.I. per acre. Do not exceed 2.72 fluid ounces per acre of **AllPro Envion rtu** which is equal to 0.007 pound permethrin A.I. per acre and 0.007 pound piperonyl butoxide A.I. per acre.

IN FLORIDA: Do not apply by aircraft unless approved by the Florida Department of Agriculture and Consumer Services.

RATES CHART						
CONDITIONS AND RATES TO USE						
ALLPRO ENVION READY-TO-USE FOR MOSQUITO CONTROL						
POUNDS A.I. PER/ACRE		APPLICATION RATES FL. OZ. PER MINUTE AT M.P.H.				FL. OZ. (ML.) FINISHED SPRAY
PERMETHRIN	PBO	5 M.P.H.	10 M.P.H.	15 M.P.H.	20 M.P.H.	PER ACRE
0.00700	0.00700	8.23	16.46	—	—	2.72 (80.44 ML.)
0.00350	0.00350	4.11	8.23	12.34	16.46	1.36 (40.22 ML.)
0.00175	0.00175	2.06	4.11	6.17	8.23	0.68 (20.11 ML.)
0.00100	0.00100	1.18	2.06	3.53	4.70	0.39 (11.53 ML.)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE AND SPILL PROCEDURES: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER HANDLING:

NONREFILLABLE CONTAINER: Do not reuse or refill this container. Triple rinse or pressure rinse container promptly after emptying. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

TRIPLE RINSE AS FOLLOWS - CONTAINERS 5 GALLONS OR LESS IN CAPACITY: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

CONTAINERS GREATER THAN 5 GALLONS IN CAPACITY: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

PRESSURE RINSE AS FOLLOWS: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure-rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

READ ENTIRE LABEL BEFORE USING

Read the entire **Directions for Use, Conditions, Disclaimer of Warranties, and Limitation of Liability** before using this product. By using this product, User and Buyer accept the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The Directions for Use of this product are believed to be adequate and should be followed carefully. However it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of **Value Garden Supply**. All such risks shall be assumed by the User or Buyer.

DISCLAIMER OF WARRANTIES: To the extent provided by law, there are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of **Value Garden Supply** is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. **Value Garden Supply** disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATION OF WARRANTIES: The exclusive remedy of the User or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability, or otherwise, shall not exceed the purchase price, or at **Value Garden Supply's** option, the replacement of the product.

EPA Reg. No 769-994
EPA Est. No. 44616-MO-1

RM# 26404-515

Manufactured by:
Value Garden Supply
P.O. Box 585
St. Joseph, MO 64502
(248) 773-7460
www.allprovector.com